

Two Patched Protein Subtypes and a Conserved Domain of Group I Proteins That Regulates Turnover*

Received for publication, August 12, 2008, and in revised form, August 29, 2008. Published, JBC Papers in Press, September 2, 2008, DOI 10.1074/jbc.M806242200

Shiho Kawamura^{‡§}, Kieran Hervold^{§1}, Felipe-Andrés Ramirez-Weber^{‡§}, and Thomas B. Kornberg^{‡2}

From the [‡]Department of Biochemistry and Biophysics, University of California, San Francisco, California 94158 and the [§]Biology Department, San Francisco State University, San Francisco, California 94132

Patched (Ptc) is a 12-cross membrane protein that binds the secreted Hedgehog protein. Its regulation of the Hedgehog signaling pathway is critical to normal development and to a number of human diseases. This report analyzes features of sequence similarity and divergence in the Ptc protein family and identifies two subtypes distinguished by novel conserved domains. We used these results to propose a rational basis for classification. We show that one of the conserved sequence regions in the C-terminal domain of Ptch1 is responsible, at least in part, for rapid turnover. This sequence is absent in the stable Ptch2 protein.

The Hedgehog (Hh) signaling pathway participates in patterning organs of both vertebrates and insects, and misregulation of the pathway has been implicated in several human diseases. Genetic and biochemical studies have identified many components that contribute to the production of active secreted Hh protein and to the receipt of and response to Hh by receiving cells. However, key features of Hh signal transduction remain poorly understood. In particular, the mechanism by which Hh engages its receptor and activates signal transduction are unclear. Current models posit that Patched (Ptc), a transmembrane protein, engages Hh on the surface of Hh-responsive cells, but how it does so and how it regulates the downstream Hh signal transduction pathway is not understood. In humans, Ptch1 has been identified as a tumor suppressor (1, 2).

Although the Ptc tertiary structure has not been determined, secondary structure predictions suggest that Ptc proteins have 12 transmembrane-spanning domains, two large extracellular loops, and a C-terminal domain (CTD).³ Structure/function studies have ascribed distinct functions to some of these domains. First, indirect evidence that Ptc binds Hh has been reported for both vertebrate Ptch1 (3–5) and invertebrate Ptc

(6–9). Based upon the phenotypes of loop 2 deletions, Hh binding has been attributed to the second extracellular loop (10). Second, five of the Ptc transmembrane segments appear to constitute a region that is conserved in the Neimann-Pick-1, hydroxymethylglutaryl-CoA reductase, and sterol regulatory element-binding protein cleavage-activating proteins (11–13) and that has been called a “sterol-sensing domain” (SSD). The NPC-1 family of proteins transports lipophilic molecules across membranes. Genetic studies suggest that Ptc negatively regulates Hh signal transduction unless bound by Hh.

Sequence analysis suggests that Ptc is structurally similar to the RND family of channels and transporters (14), and evidence has been reported for Hh-sensitive repression of signal transduction being effected by a transporter function of Ptc either for sterols (15, 16) or for vitamin D₃ (17). Transporter activity would be consistent with the lack-of-function phenotype of Ptc protein mutants with altered residues that are conserved in and are required for activity of bacterial RND transporters (9) or of NPC-1 (18, 19). One RND family transporter, the proton-driven *Escherichia coli* AcrB protein, is a homotrimer with a central pore (20), and recent evidence suggests that Ptc is also a trimer (8).

A number of distinct functions are also encoded in the Ptc CTD. Activation of the Hh pathway correlates with relocalization of Ptc from the plasma membrane to endosomes (21–24), movement that is dependent upon the CTD. The *ptc*^{I3} mutant has a missense change in the CTD (E1172K) and a partial loss-of-function phenotype; its endosomal localization is decreased (19). Deletion of the CTD also affects localization and results in plasma membrane accumulation (8, 24). The CTD can form a homotrimer, and it contains a PPAY sequence, a putative target site for the Nedd4 ubiquitin ligase (8). CTD deletions are more stable than the wild-type protein (8, 9). Several mutations in the CTD of human Ptch1 have been identified in human cancers (25).

Drosophila has a single *ptc* gene, but most vertebrates have two, customarily named Patched1 (*Ptch1*) and Patched2 (*Ptch2*). In general, these gene designations reflect the relative order of discovery, not function or sequence subtype. This study describes a bioinformatics analysis with the goal of establishing a basis for categorizing the members of the Ptc protein family. Because the CTD sequences are significantly less well conserved than other Ptc domains (8), their variability was used to define two subtypes. One (Group I) includes the invertebrate Ptc proteins and the Ptch1 proteins of human, mouse, dog, frog, chick, zebrafish, monkey, and rat as well as the Ptch2 protein of zebrafish. Group I CTDs are characterized by having two con-

* This work was supported, in whole or in part, by National Institutes of Health Grant GM077407 (to T. B. K.). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

[†] Deceased November 1, 2007.

¹ Present address: Xoma, 2910 Seventh St., Berkeley, CA 94710.

² To whom correspondence should be addressed: Dept. of Biochemistry and Biophysics, University of California, 1550 4th St., San Francisco, CA 94158. Fax: 415-476-3892; E-mail: tkornberg@ucsf.edu.

³ The abbreviations used are: CTD, C-terminal domain; SSD, sterol-sensing domain; RLM-RACE, RNA ligase-mediated rapid amplification of cDNA ends.

served domains, a PPXY sequence, and a size between 205 and 279 residues. The CTDs of Group II proteins have abbreviated versions of the two conserved domains, lack a PPXY sequence, and are shorter, containing between 75 and 109 residues; they likely represent a novel mammalian divergence. We also show that the PPXY sequence in the C-terminal cytoplasmic tail of *Ptch1* is responsible, at least in part, for rapid turnover. The *Ptch2* protein is relatively stable; it lacks a PPXY sequence.

EXPERIMENTAL PROCEDURES

Gene Predictions—The amino acid sequences of *Ptc* orthologs were collected from the GenBankTM Data Bank. To expand the set of *Ptc* CTD sequences, genomic sequences were also accessed using the UCSC Genome Browser (26) and Fly-Base. Using these sources and tools, we gathered the genomic sequences of putative *Ptc* orthologs based upon their homology to *ptc* genes in human, mouse, and *Drosophila melanogaster*.

Homologous genomic regions, plus 3 kb of additional sequence on either end, were analyzed with three gene prediction programs: Genescan (27), Augustus (28), and GeneID (29). Results from the programs varied slightly, and choices between different gene predictions were made as follows. The predictions were aligned with the sequences of “evolutionary relatives”: *D. melanogaster* *Ptc* for insects and mouse and human *Ptc1* and *Ptc2* for vertebrates. Alignments were created using ClustalW, tCOFFEE (30), or MUSCLE Version 3.7 (31). Results were displayed in Jalview (32). Because introns were more frequently mistaken for exons than the reverse and because we are concerned primarily with the CTD, predictions were chosen to minimize false inserts within the tail region.

mRNA Extraction and cDNA Formation—Seven *Drosophila* species were analyzed: *D. ananassae*, *D. mojavensis*, *D. persimilis*, *D. pseudoobscura*, *D. virilis*, *D. simulans*, and *D. yakuba*. RNA was extracted from 15 third-instar larvae and fractionated by selection with oligo(dT), and first-strand cDNA was synthesized using the RETROscript[®] kit (Ambion). Mid-neurula stage *Xenopus tropicalis* embryos (stages 14–18) were from R. Harland (University of California, Berkeley, CA); zebrafish embryos (18-somite stage) were from S. Amacher (University of California, Berkeley). mRNA was purified with the RNeasy minikit (Qiagen Inc.). cDNAs were synthesized using ~2.0 µg of total mRNA and the FirstChoice[®] RLM-RACE kit (Ambion). *Canis familiaris* and Rhesus monkey (*Macaca mulatta*) cDNAs were purchased from BioChain Institute.

Primer Design for Reverse Transcription-PCR—Primers were designed based upon predicted CTD sequences and were chosen to span ~300 bp of intron. PCR-amplified cDNA was purified after separation by agarose gel electrophoresis. Primer sequences were as follows: *D. ananassae*, GTCAAGTTCGAGGGCTTT and ACTCGTAAAGTTGTAGCTGCG (Rev1); *D. simulans*, CCTGCCCAACTATCCATC and CTCTTGCTGCTGCGCAC; *D. mojavensis* and *D. virilis*, CAAGACACTGATTGGACACATAC and Rev1; *D. pseudoobscura*, CGACCTGAAGATACCCAAGA and Rev1; *D. yakuba*, CTCAGATGCCCTTTTACCTTC and Rev1; *C. familiaris*, TGCTGGGAGTGCTGATG and GTTGGAGCTACCGCCC; *M. mulatta*, GTTTTGCTTCCCGTGCTTT and GTTGGAGCTGCTTC-

CCC; *Danio rerio*, TGCTATTCCTGTGGTCATCCT (first RLM-RACE forward primer) and TGCCTCCTCCTATGAA-CCAC and CTATTTACCTAGCATCTGGTTAGCTTG (PCR primers); and *X. tropicalis* TTTGCTGTCTGGCAATTTT (first RLM-RACE forward primer) and ATGATTCCCAGCATGTTC and TTATTTGAGGCATTTCCCTGT (PCR primers).

Amplifications were performed using the High Fidelity PCR Master (Roche Applied Science). For sequencing, PCR-amplified *ptc* fragments were cloned into the pCRII-topo vector (Invitrogen). Six individual colonies of each species were sequenced.

Expression Constructs—Mouse *Ptch1* and *Ptch2* were cloned into pAcGFP-C3 (Clontech), generating fusion proteins with an N-terminal green fluorescent protein: *Ptch1*^{C2}, *Ptch1*-(1–1162)/*Ptch2*-(1113–1171); *Ptch2*^{C1}, *Ptch2*-(1–1112)/*Ptch1*-(1163–1377); *Ptch1*^{ΔC}, *Ptch1*-(1–1162); and *Ptch1*^{ΔA}, PPPY (positions 1299–1302) to AAAA.

Western Analysis—NIH 3T3 cells were cultured in high-glucose Dulbecco's modified Eagle's medium with 10% (v/v) bovine calf serum. Cells (40,000 cells/cm²) were seeded in 6-well plates, incubated for 18 h (to 90–95% confluence), and transfected with Lipofectamine 2000 (Invitrogen). After 48 h, cycloheximide (100 µM; Sigma) was added. After the indicated periods, cells were rinsed with phosphate-buffered saline; detached with 500 µl of 0.25% trypsin/EDTA; and after the addition of 500 µl of medium, collected by centrifugation at 7000 rpm for 5 min. Cells were rinsed with 500 µl of cold phosphate-buffered saline and lysed with 100 µl of cold lysis buffer (0.5% Nonidet P-40, 50 mM NaCl, 50 mM Tris (pH 7.5) and Complete protease inhibitor mixture (Roche Applied Science)) by pipetting ~30 times. After 30 min of agitation on a rocker at 4 °C, extracts were centrifuged at maximum speed for 30 min, and 20 µl of sample was mixed with 5 µl of 5× Laemmli sample buffer and incubated for 15 min at 4 °C. Twenty µl of each sample was loaded onto a 6% SDS-polyacrylamide gel (2% polyacrylamide stacking gel) that had been pre-run for 20 min at 70 V.

Western blots were prepared on polyvinylidene difluoride membranes by semidry electrotransfer, blocked with 5% dry milk in TBST (0.08% Tween 20, 150 mM NaCl, and 20 mM Tris) for 30 min at room temperature, and incubated overnight at 4 °C with rabbit anti-green fluorescent protein antibody (1:1000 in 5% dry milk/TBST). After washing three times with 5% dry milk/TBST for 1 h, membranes were incubated for 1.5 h with horseradish peroxidase-conjugated goat anti-rabbit IgG (diluted 1:1000 in 5% dry milk/TBST). Membranes were washed twice with 5% dry milk/TBST for 30 min and rinsed with TBST for 10 min. Detection was with the ECL PlusTM Western blotting detection system (Amersham Biosciences). Blots were quantified with NIH Image J. The intensity of each construct was compared with the tubulin loading control, and the ratio at 0 h was set at 100%.

RESULTS

Conservation of *Ptc* Domains—Conservation of *Ptc* protein sequences during vertebrate and invertebrate evolution has been significant. To investigate how different regions of *Ptc*

Regulation of *Ptch1* and *Ptch2* Turnover

have either changed or been conserved, we assembled a group of invertebrate and vertebrate Ptc sequences for comparison. Thirteen sequences of vertebrate Ptc proteins were obtained from the NCBI Database (see Fig. 2). In addition, we prepared and partially sequenced *Ptch* cDNAs from *C. familiaris*, *M. musculus*, and *X. tropicalis*; the total number of CTD sequences in our data set was 16. Thirteen sequences of invertebrate Ptc proteins were obtained from the NCBI and FlyBase Databases. We verified the Ptc CTD sequences for six of the *Drosophila* species (see Fig. 2).

To estimate the general similarities of Ptc proteins, we first compared the sequences of *D. melanogaster* Ptc, which was the first Ptc protein identified, and human and mouse *Ptch1* and *Ptch2*. Pairwise ClustalW comparisons of these proteins revealed the near identity of human and mouse *Ptch1* proteins and of human and mouse *Ptch2* proteins and the high degree of similarity (>50%) of these five proteins (Table 1). These alignments also revealed that the general topology of these proteins is similar, a result that is consistent with the similar distribution of predicted membrane-spanning domains in each protein (data not shown).

We extended this analysis to the entire data set and also focused on four domains: the predicted extracellular loops 1 and 2, the SSD, and the CTD. Pairwise comparisons of the full-

length proteins and of four domains were made for both the invertebrate and vertebrate sequence data sets. These analyses revealed sequence similarities of >50% for the full-length proteins and >65% in both extracellular loops and the SSD. Variation in the CTDs was significantly higher (Fig. 1).

Vertebrate *Ptch* Proteins Have Either of Two CTD Subtypes—Further analysis of the vertebrate CTDs with ClustalW revealed two highly conserved subtypes that share >50% homology with each other (Table 2). The first group (Group I) includes dog, human, monkey, mouse, rat, and *Xenopus laevis* *Ptch1* and *X. tropicalis* Ptc. Group II includes cow, dog, human, mouse, and rat *Ptch2*. The CTD sequences of *X. laevis* *Ptch2* and zebrafish *Ptch1* and *Ptch2* have low homology to either group. Although we cannot evaluate why these sequences are so highly divergent (whether the differences are real or an artifact of isolation and cloning), as described below, the conservation they do have suggests that they should be included among the Group I orthologs.

Two other features individuate these two groups (Table 3). First, Group I CTDs are significantly longer than Group II CTDs. With the exception of the two fish CTDs, all Group I CTDs exceed 205 residues. In contrast, none of the Group II CTDs is longer than 109 residues. Second, only the Group I CTDs include the PPXY sequence. In the *D. melanogaster* CTD, the PPXY residues have been implicated in the regulated turnover of the Ptc protein (8), and mutations in the PPXY sequence increase the protein half-life. CTD deletion mutants of both the *D. melanogaster* and mouse Ptc1 proteins also have reduced rates of degradation (Fig. 4) (8). The PPXY motif is not present in any of the Group II CTDs.

Conserved Sequences in the *Ptch* CTD—Phylogenetic relationships predicted by the relative degree of sequence similarity between Ptc proteins are consistent with a model in which Group I and II proteins represent distinct subtypes (Fig. 2). We generated phylogenetic trees by the method of “neighbor joining” based on BLOSUM62 for all of the Ptc proteins in our data set as well as for each of the four domains. Similar results were obtained for all five trees: the invertebrate Group I and II sequences were placed on separate branches in each calculation.

Sequence alignment performed by ClustalW, MUSCLE, and tCOFFEE identified two islands of significant conservation in the vertebrate Group I and II CTDs as well as the invertebrate CTDs. For purposes of illustration (Fig. 3), we used ClustalW to align vertebrate Group I and II CTDs and extracted a CTD “consensus” sequence for each type. We also generated an invertebrate CTD consensus sequence by aligning the *D. melanogaster*, *Aedes aegypti*, and *Anopheles gambiae*

TABLE 1
Similarity analysis of Ptc proteins

The calculated values of the percent similarity between full-length human (*Homo sapiens*), mouse (*Mus musculus*), and *D. melanogaster* Ptc proteins are given below.

	<i>H. sapiens</i> 1	<i>H. sapiens</i> 2	<i>M. musculus</i> 1	<i>M. musculus</i> 2
<i>H. sapiens</i> 2	68			
<i>M. musculus</i> 1	97	61		
<i>M. musculus</i> 2	60	94	60	
<i>D. melanogaster</i>	51	51	51	51




FIGURE 1. Sequence variance in vertebrate and invertebrate Ptc proteins. The sequence similarity value of pairwise ClustalW alignments is plotted for the full-length proteins and for loop 1, the SSD, loop 2, and the CTD. Each circle represents a percent similarity value calculated for one of the pairwise comparisons among invertebrate (left) and vertebrate (right) proteins.

TABLE 2
Pairwise comparisons of vertebrate Ptch sequences

The percent values for sequence similarity between the CTDs in each Ptch protein in the data set are given below; values $\geq 50\%$ are indicated in boldface.

	<i>C. familiaris 1</i>	<i>D. rerio 1</i>	<i>D. rerio 2</i>	<i>Gallus gallus 1</i>	<i>H. sapiens 1</i>	<i>M. mulatta</i>	<i>M. musculus 1</i>	<i>Rattus norvegicus 1</i>	<i>X. laevis 1</i>	<i>X. laevis 2</i>	<i>X. tropicalis</i>	<i>Bos taurus 2</i>	<i>C. familiaris 2</i>	<i>H. sapiens 2</i>	<i>M. musculus 2</i>	
<i>D. rerio 1</i>	11															
<i>D. rerio 2</i>	17	31														
<i>G. gallus 1</i>	75	11	18													
<i>H. sapiens 1</i>	90	11	16	77												
<i>M. mulatta</i>	91	11	16	78	97											
<i>M. musculus 1</i>	90	11	17	79	94	95										
<i>R. norvegicus 1</i>	90	11	17	79	94	95	100									
<i>X. laevis 1</i>	63	10	16	68	63	65	65	65								
<i>X. laevis 2</i>	36	16	12	37	37	38	37	37	37							
<i>X. tropicalis</i>	49	2	3	53	48	50	50	50	67	29						
<i>B. taurus 2</i>	13	17	13	13	13	13	13	13	13	13	8					
<i>C. familiaris 2</i>	11	12	9	11	12	12	12	12	10	11	8	76				
<i>H. sapiens 2</i>	13	19	15	13	13	13	14	14	14	13	8	85	73			
<i>M. musculus 2</i>	13	22	17	12	12	13	13	13	11	12	7	66	56	67		
<i>R. norvegicus 2</i>	13	22	17	12	13	13	13	13	12	12	7	68	58	70	95	

TABLE 3

Characteristics of Group I and II CTDs

Listed are the number of residues in each CTD and the residue location of the PPXY sequence that is present in 9 of the 11 Group I proteins.

	Total length	PPXY
Group I		
<i>C. familiaris 1</i>	281	142
<i>D. rerio 1</i>	60	
<i>D. rerio 2</i>	58	
<i>G. gallus 1</i>	272	139
<i>H. sapiens 1</i>	276	142
<i>M. mulatta</i>	279	142
<i>M. musculus</i>	277	142
<i>R. norvegicus 1</i>	277	142
<i>X. laevis 1</i>	250	126
<i>X. laevis 2</i>	262	176
<i>X. tropicalis</i>	205	72
Group II		
<i>B. taurus 2</i>	100	
<i>C. familiaris 2</i>	109	
<i>H. sapiens 2</i>	96	
<i>M. musculus 2</i>	75	
<i>R. norvegicus 2</i>	75	

sequences. Fig. 3 presents an alignment of this “invertebrate CTD” consensus sequence with the consensus sequences of the vertebrate Group I and II CTDs. Two features of these alignments are noteworthy. First, the region of 30–40 residues at the immediate N terminus of the CTD is revealed as a domain of conservation in all three consensus sequences. We designated this region “Domain I.” Its conservation in the three sequences is $>67\%$ within the N-terminal 15 residues, and whereas the Group II homology does not extend further, the region of $>50\%$ conservation extends to 27 residues in the invertebrate Group I domains. Domain I of the zebrafish CTDs is the longer variety, consistent with a Group I designation. Second, “Domain II,” a more C-terminal region of conservation, extends for ~ 21 residues and is also present in all three CTD types.

Role of the CTD in Regulating Turnover of Ptc1 and Ptc2—The CTD of the *D. melanogaster* Ptc protein regulates both turnover and internalization (8). The wild-type protein is unstable, with a half-life of <2 h. Its CTD contains a potential target sequence (PPAY) for the Nedd4 ubiquitin ligase, and mutant protein that lacks either the C-terminal half of the CTD or the PPAY sequence is significantly more stable (8). Because our comparative sequence analysis (above) revealed that Group I Ptc proteins have a potential neddylation sequence (PPPY) but Group II proteins do not, we investigated whether the PPPY sequence in vertebrate Ptc proteins has a role in regulating turnover.

As shown in Fig. 4, Ptch1 was unstable ($t_{1/2} = 1.7$ h), whereas Ptch2 was stable ($t_{1/2} > 18$ h). Deleting the CTD of Ptch1 stabilized the protein ($t_{1/2} > 17$ h); swapping the CTDs of Ptch1 and Ptch2 stabilized Ptch1 ($t_{1/2} = 5.8$ h) and destabilized Ptch2 ($t_{1/2} = 6.2$ h). These properties indicate that the CTDs of Group I and II proteins help to determine lifetime. Mutating the PPPY sequence in the Ptch1 CTD to AAAA stabilized Ptch1 ($t_{1/2} > 9$ h). We conclude that the CTD of the vertebrate Ptc proteins plays a major role in regulating stability and that the PPPY sequence is an essential component of the mechanism that controls Ptc1 turnover.

Regulation of *Ptch1* and *Ptch2* Turnover

DISCUSSION

The Ptc protein family is characterized by shared topology and high sequence conservation. All members are predicted to have 12 membrane-spanning domains, an SSD, two large extracellular loops, and a C-terminal cytoplasmic tail. Several lines of evidence suggest that two subtypes of Ptc protein compose this family. This evidence includes predicted phylogenetic relationships among the family members as well as separate comparisons of sequence similarities within extracellular loops 1 and 2, the SSD, and the CTD. Most Ptc proteins, including the invertebrate orthologs, are of the Group I type. Although the known family members represent products from the small and selected set of genomes that were available to us, it bears noting that Group II orthologs have been identified only in mammals. It follows that the Group II subtype may be unique to mammals and may represent a divergence in that lineage. Whereas it is clear that Group I proteins function as tumor suppressors and as key negative regulators of Hedgehog signal transduction, the

function of the Group II proteins has not been elucidated. Studies of mouse *Ptch2* mutants suggest that this Group II protein modulates tumorigenesis associated with *Ptch1* haploinsufficiency (33), but how *Ptch2* affects *Ptch2* activity or downstream functions is not known.

Although sequence variance in the CTD is significantly greater than in other domains, the islands of conservation in it may represent regions with conserved function. We designated these regions Domain I, Domain II, and PPXY, and available genetic evidence suggests that all three are functionally important. Mutations in the *D. melanogaster* PPXY sequence increase the protein half-life (8), suggesting that this sequence motif functions to limit the lifetime of Ptc protein. In this work, we have shown that the PPPY sequence in mouse *Ptch1* has a similar function. It is interesting to speculate that Group II proteins may not be regulated by the same mechanism and that the absence of the PPXY motif might indicate a different role for this protein. Domain I may also be critical to *Ptch* function. Among 12 CTD mutations that are listed in the PTCH Mutation Database of human mutations (25), three (T1195I, P1196S, and S1197F) alter conserved residues in Domain I. These three mutations were identified in patients with basal cell carcinoma, suggesting that these mutant proteins may not respond to Hh or its absence in a normal way. Presumably, the pathway is abnormally active in these mutant backgrounds, and given the negative role of Ptc, these mutations might decrease Ptc activity either by reducing the half-life of the protein or by affecting its localization. Although Domain II mutations in human Ptc have not been reported, studies of the *D. melanogaster* CTD with mutations in Domain II have revealed that Domain II is essential for self-association of the CTD (8).




FIGURE 2. **Phylogenetic tree of Ptc orthologs.** The phylogenetic tree for the data set of vertebrate and invertebrate proteins was calculated using MUSCLE Version 3.7 (31) and Jalview Version 12.0.0 with default settings for alignment matrix and other parameters.




FIGURE 3. **Alignment of vertebrate Group I and II and invertebrate CTD sequences.** Group I, Group II, and invertebrate CTD sequences were aligned using ClustalW and are displayed in Jalview, a multiple alignment editor that identifies conserved residues (blue, top rows); assesses "quality" of conservation (middle rows) by analyzing the proportion of identical conservative changes and non-conservation (scaled high (yellow), intermediate (brown), and low (black)); and defines an overall "consensus" (bottom rows). Islands of high sequence conservation (Domains I and II) and the PPPY sequence are indicated by boxes.




FIGURE 4. The CTD regulates turnover of Ptch proteins. The stability of Ptch proteins in transfected NIH 3T3 cells was assessed by Western analysis of cells treated with cycloheximide (CHX) for the indicated times. Wild-type mouse Ptch1 and Ptch2, Ptch1 substituted with a Ptch2 CTD (Ptch1^{C2}), Ptch2 substituted with a Ptch1 CTD (Ptch2^{C1}), Ptch1 deleted of its CTD (Ptch1^{ΔC}), and Ptch1 with the PPPY sequence in its CTD changed to AAAA (Ptch1^{4A}) are compared. Red, Ptch1 residues; blue, Ptch2 residues; black, mutated PPPY in CTD.

It remains for future work to investigate whether the Domain II sequences in vertebrate Group I and II proteins contribute a similar function.

Acknowledgments—We thank members of the Kornberg laboratory and Patricia Babbitt for technical help and comments on the manuscript, Michiel Bagnat for technical advice, and Bruce Macher, Laura Burrus, Ravinder Sehgal, and Diana Chu (San Francisco State University) for guidance and support.

REFERENCES

- Hahn, H., Wicking, C., Zaphiropoulos, P. G., Gailani, M. R., Shanley, S., Chidambaram, A., Vorechovsky, I., Holmberg, E., Uden, A. B., Gillies, S., Negus, K., Smyth, I., Pressman, C., Leffell, D. J., Gerrard, B., Goldstein, A. M., Dean, M., Toftgard, R., Chenevix-Trench, G., Wainwright, B., and Bale, A. E. (1996) *Cell* **85**, 841–851
- Johnson, R. L., Rothman, A. L., Xie, J., Goodrich, L. V., Bare, J. W., Bonifas, J. M., Quinn, A. G., Myers, R. M., Cox, D. R., Epstein, E. H., Jr., and Scott, M. P. (1996) *Science* **272**, 1668–1671
- Carpenter, D., Stone, D. M., Brush, J., Ryan, A., Armanini, M., Frantz, G., Rosenthal, A., and de Sauvage, F. J. (1998) *Proc. Natl. Acad. Sci. U. S. A.* **95**, 13630–13634
- Marigo, V., Davey, R. A., Zuo, Y., Cunningham, J. M., and Tabin, C. J. (1996) *Nature* **384**, 176–179
- Stone, D. M., Hynes, M., Armanini, M., Swanson, T. A., Gu, Q., Johnson, R. L., Scott, M. P., Pennica, D., Goddard, A., Phillips, H., Noll, M., Hooper, J. E., de Sauvage, F., and Rosenthal, A. (1996) *Nature* **384**, 129–134
- Fuse, N., Maiti, T., Wang, B., Porter, J. A., Hall, T. M., Leahy, D. J., and Beachy, P. A. (1999) *Proc. Natl. Acad. Sci. U. S. A.* **96**, 10992–10999
- Ingham, P. W., Taylor, A. M., and Nakano, Y. (1991) *Nature* **353**, 184–187
- Lu, X., Liu, S., and Kornberg, T. B. (2006) *Genes Dev.* **20**, 2539–2551
- Taipale, J., Cooper, M. K., Maiti, T., and Beachy, P. A. (2002) *Nature* **418**, 892–897
- Briscoe, J., Chen, Y., Jessell, T. M., and Struhl, G. (2001) *Mol. Cell* **7**, 1279–1291
- Carstea, E. D., Morris, J. A., Coleman, K. G., Loftus, S. K., Zhang, D., Cummings, C., Gu, J., Rosenfeld, M. A., Pavan, W. J., Krizman, D. B., Nagle, J., Polymeropoulos, M. H., Sturley, S. L., Ioannou, Y. A., Higgins, M. E., Comly, M., Cooney, A., Brown, A., Kaneski, C. R., Blanchette-Mackie, E. J., Dwyer, N. K., Neufeld, E. B., Chang, T. Y., Liscum, L., Strauss, J. F., III, Ohno, K., Zeigler, M., Carmi, R., Sokol, J., Markie, D., O'Neill, R. R., van Diggelen, O. P., Elleder, M., Patterson, M. C., Brady, R. O., Vanier, M. T., Pentchev, P. G., and Tagle, D. A. (1997) *Science* **277**, 228–231
- Hua, X., Nohturfft, A., Goldstein, J. L., and Brown, M. S. (1996) *Cell* **87**, 415–426
- Osborne, T. F., and Rosenfeld, J. M. (1998) *Curr. Opin. Lipidol.* **9**, 137–140
- Tseng, T. T., Gratwick, K. S., Kollman, J., Park, D., Nies, D. H., Goffeau, A., and Saier, M. H., Jr. (1999) *J. Mol. Microbiol. Biotechnol.* **1**, 107–125
- Corcoran, R. B., and Scott, M. P. (2006) *Proc. Natl. Acad. Sci. U. S. A.* **103**, 8408–8413
- Dwyer, J. R., Sever, N., Carlson, M., Nelson, S. F., Beachy, P. A., and Parhami, F. (2007) *J. Biol. Chem.* **282**, 8959–8968
- Bijlsma, M. F., Spek, C. A., Zivkovic, D., van de Water, S., Rezaee, F., and Peppelenbosch, M. P. (2006) *PLoS Biol.* **4**, e232
- Martin, V., Carrillo, G., Torroja, C., and Guerrero, I. (2001) *Curr. Biol.* **11**, 601–607
- Strutt, H., Thomas, C., Nakano, Y., Stark, D., Neave, B., Taylor, A. M., and Ingham, P. W. (2001) *Curr. Biol.* **11**, 608–613
- Murakami, S., Nakashima, R., Yamashita, E., and Yamaguchi, A. (2002) *Nature* **419**, 587–593
- Capdevila, J., Pariente, F., Sampedro, J., Alonso, J. L., and Guerrero, I. (1994) *Development (Camb.)* **120**, 987–998
- Denef, N., Neubuser, D., Perez, L., and Cohen, S. M. (2000) *Cell* **102**, 521–531
- Incardona, J. P., Gruenberg, J., and Roelink, H. (2002) *Curr. Biol.* **12**, 983–995
- Zhu, A. J., Zheng, L., Suyama, K., and Scott, M. P. (2003) *Genes Dev.* **17**, 1240–1252
- Lindstrom, E., Shimokawa, T., Toftgard, R., and Zaphiropoulos, P. G. (2006) *Hum. Mutat.* **27**, 215–219
- Kent, W. J., Baertsch, R., Hinrichs, A., Miller, W., and Haussler, D. (2003) *Proc. Natl. Acad. Sci. U. S. A.* **100**, 11484–11489
- Burge, C., and Karlin, S. (1997) *J. Mol. Biol.* **268**, 78–94
- Stanke, M., Schoffmann, O., Morgenstern, B., and Waack, S. (2006) *BMC Bioinformatics* **7**, 62
- Parra, G., Blanco, E., and Guigo, R. (2000) *Genome Res.* **10**, 511–515
- Notredame, C., Higgins, D. G., and Heringa, J. (2000) *J. Mol. Biol.* **302**, 205–217
- Edgar, R. C. (2004) *Nucleic Acids Res.* **32**, 1792–1797
- Clamp, M., Cuff, J., Searle, S. M., and Barton, G. J. (2004) *Bioinformatics (Oxf.)* **20**, 426–427
- Lee, Y., Miller, H. L., Russell, H. R., Boyd, K., Curran, T., and McKinnon, P. J. (2006) *Cancer Res.* **66**, 6964–6971